

Dual algebras and A-measures.

Marek Kosiek

Joint work with Krzysztof Rudol

A - an arbitrary function algebra

A - an arbitrary function algebra

The main subject of our investigation:

Properties of the spectrum of A^{**} .

A - an arbitrary function algebra

The main subject of our investigation:

Properties of the spectrum of A^{**} .

Motivation:

A - an arbitrary function algebra

The main subject of our investigation:

Properties of the spectrum of A^{**} .

Motivation:

- A - measures problem

A - an arbitrary function algebra

The main subject of our investigation:

Properties of the spectrum of A^{**} .

Motivation:

- A - measures problem
- the problem for which $G \subset \mathbb{C}^n$ the algebra $H^\infty(G)$ is a dual algebra

A - an arbitrary function algebra

The main subject of our investigation:

Properties of the spectrum of A^{**} .

Motivation:

- A - measures problem
- the problem for which $G \subset \mathbb{C}^n$ the algebra $H^\infty(G)$ is a dual algebra
- the application of dual algebras in functional calculus for bounded operators in Hilbert spaces

A - an arbitrary function algebra

The main subject of our investigation:

Properties of the spectrum of A^{**} .

Motivation:

- A - measures problem
- the problem for which $G \subset \mathbb{C}^n$ the algebra $H^\infty(G)$ is a dual algebra
- the application of dual algebras in functional calculus for bounded operators in Hilbert spaces
- connections with the Corona problem

As an application of our main result we have obtained:

As an application of our main result we have obtained:

- a general positive solution for A - measures problem

As an application of our main result we have obtained:

- a general positive solution for A - measures problem
- the duality of $H^\infty(G)$ algebra for some classes of bounded domains $G \subset \mathbb{C}^n$

Definition

A is a function algebra on a compact set X iff $A \subset C(X)$,
 A contains constants and separates the points of X

Definition

A is a function algebra on a compact set X iff $A \subset C(X)$,
 A contains constants and separates the points of X

Let $\phi, \psi \in \sigma(A)$

$$\phi \sim \psi \stackrel{\text{df}}{\iff} \|\phi - \psi\| < 2$$

Definition

A is a function algebra on a compact set X iff $A \subset C(X)$,
 A contains constants and separates the points of X

Let $\phi, \psi \in \sigma(A)$

$$\phi \sim \psi \stackrel{\text{df}}{\iff} \|\phi - \psi\| < 2$$

Definition

The equivalence classes in the above equivalence relation are called **Gleason parts** of A .

Definition

A is a function algebra on a compact set X iff $A \subset C(X)$,
 A contains constants and separates the points of X

Let $\phi, \psi \in \sigma(A)$

$$\phi \sim \psi \stackrel{\text{df}}{\iff} \|\phi - \psi\| < 2$$

Definition

The equivalence classes in the above equivalence relation are called **Gleason parts** of A .

We assume $\sigma(A) = X$.

- $\mathcal{M} \subset M(X) := C(X)^*$ is a band if it is a closed subspace and $\mu \in \mathcal{M}, \nu \ll |\mu| \implies \nu \in \mathcal{M}$

- $\mathcal{M} \subset M(X) := C(X)^*$ is a band if it is a closed subspace and $\mu \in \mathcal{M}, \nu \ll |\mu| \implies \nu \in \mathcal{M}$
- every measure $\mu \in M(X)$ has a unique Lebesgue decomposition $\mu = \mu_{\mathcal{M}} + \mu_s$ where $\mu_{\mathcal{M}} \in \mathcal{M}$ and μ_s is singular to each measure in \mathcal{M}

- $\mathcal{M} \subset M(X) := C(X)^*$ is a band if it is a closed subspace and $\mu \in \mathcal{M}, \nu \ll |\mu| \implies \nu \in \mathcal{M}$
- every measure $\mu \in M(X)$ has a unique Lebesgue decomposition $\mu = \mu_{\mathcal{M}} + \mu_s$ where $\mu_{\mathcal{M}} \in \mathcal{M}$ and μ_s is singular to each measure in \mathcal{M}
- \mathcal{M} is a reducing band (with respect to A) if $\mu \in A^\perp \implies \mu_{\mathcal{M}} \in A^\perp$

- $\mathcal{M} \subset M(X) := C(X)^*$ is a band if it is a closed subspace and $\mu \in \mathcal{M}, \nu \ll |\mu| \implies \nu \in \mathcal{M}$
- every measure $\mu \in M(X)$ has a unique Lebesgue decomposition $\mu = \mu_{\mathcal{M}} + \mu_s$ where $\mu_{\mathcal{M}} \in \mathcal{M}$ and μ_s is singular to each measure in \mathcal{M}
- \mathcal{M} is a reducing band (with respect to A) if $\mu \in A^\perp \implies \mu_{\mathcal{M}} \in A^\perp$
- ν is a representing measure for $x \in X = \sigma(A)$ if $f(x) = \int f d\nu$ for $f \in A$

- $\mathcal{M} \subset M(X) := C(X)^*$ is a band if it is a closed subspace and $\mu \in \mathcal{M}, \nu \ll |\mu| \implies \nu \in \mathcal{M}$
- every measure $\mu \in M(X)$ has a unique Lebesgue decomposition $\mu = \mu_{\mathcal{M}} + \mu_s$ where $\mu_{\mathcal{M}} \in \mathcal{M}$ and μ_s is singular to each measure in \mathcal{M}
- \mathcal{M} is a reducing band (with respect to A) if $\mu \in A^\perp \implies \mu_{\mathcal{M}} \in A^\perp$
- ν is a representing measure for $x \in X = \sigma(A)$ if $f(x) = \int f d\nu$ for $f \in A$
- for $G \subset X$ we denote by \mathcal{M}_G the band generated by G i.e. the smallest band containing all measures representing for points in G

- $\mathcal{M} \subset M(X) := C(X)^*$ is a band if it is a closed subspace and $\mu \in \mathcal{M}, \nu \ll |\mu| \implies \nu \in \mathcal{M}$
- every measure $\mu \in M(X)$ has a unique Lebesgue decomposition $\mu = \mu_{\mathcal{M}} + \mu_s$ where $\mu_{\mathcal{M}} \in \mathcal{M}$ and μ_s is singular to each measure in \mathcal{M}
- \mathcal{M} is a reducing band (with respect to A) if $\mu \in A^\perp \implies \mu_{\mathcal{M}} \in A^\perp$
- ν is a representing measure for $x \in X = \sigma(A)$ if $f(x) = \int f d\nu$ for $f \in A$
- for $G \subset X$ we denote by \mathcal{M}_G the band generated by G i.e. the smallest band containing all measures representing for points in G
- if G is a Gleason part then \mathcal{M}_G is a reducing band

- $(C(X)^*)^* = M(X)^* \approx C(Y)$ for some hyperstonean compact space Y

- $(C(X)^*)^* = M(X)^* \approx C(Y)$ for some hyperstonean compact space Y
- each $f \in C(X)$ can be treated as a functional on $M(X)$ and consequently as an element of $C(Y)$ by the formula

$$\langle f, \mu \rangle = \int f d\mu \quad \text{for } \mu \in M(X)$$

- $(C(X)^*)^* = M(X)^* \approx C(Y)$ for some hyperstonean compact space Y
- each $f \in C(X)$ can be treated as a functional on $M(X)$ and consequently as an element of $C(Y)$ by the formula

$$\langle f, \mu \rangle = \int f d\mu \quad \text{for } \mu \in M(X)$$

- for $F \in C(Y)$, $\mu \in M(X)$ there is a unique measure $\tilde{\mu} \in M(Y) := C(Y)^*$ such that $\langle F, \mu \rangle = \int F d\tilde{\mu}$

Theorem

If G is a Gleason part of A then the weak-star closure \overline{G}^{ws} of G is a closed-open subset of Y . Moreover

$$Y \setminus \overline{G}^{ws} = \overline{X \setminus G}^{ws}, \quad (\overline{\mathcal{M}_G}^{ws})^s = \overline{(\mathcal{M}_G^s)}^{ws}, \quad \overline{\mathcal{M}_G}^{ws} = M(\overline{G}^{ws}),$$

and $\overline{\mathcal{M}_G}^{ws}$ is a reducing band for A^{**} .

Theorem

If G is a Gleason part of A then the weak-star closure \overline{G}^{ws} of G is a closed-open subset of Y . Moreover

$$Y \setminus \overline{G}^{ws} = \overline{X \setminus G}^{ws}, \quad (\overline{\mathcal{M}_G}^{ws})^s = \overline{(\mathcal{M}_G^s)}^{ws}, \quad \overline{\mathcal{M}_G}^{ws} = M(\overline{G}^{ws}),$$

and $\overline{\mathcal{M}_G}^{ws}$ is a reducing band for A^{**} .

Corollary

There exists a characteristic function $F_0 \in A^{**}$ vanishing exactly on $Y \setminus \overline{G}^{ws}$ and the projection associated with the decomposition $M(Y) = \overline{\mathcal{M}_G}^{ws} + \overline{\mathcal{M}_G^s}^{ws}$ is exactly the multiplication by F_0 .

- G - a Gleason part of A

- G - a Gleason part of A
- $H^\infty(\mathcal{M}_G)$ - the weak-star closure of A in \mathcal{M}_G^*

- G - a Gleason part of A
- $H^\infty(\mathcal{M}_G)$ - the weak-star closure of A in \mathcal{M}_G^*
- by the definition of $H^\infty(\mathcal{M}_G)$, the values of its every element are uniquely defined on each $x \in G$

- G - a Gleason part of A
- $H^\infty(\mathcal{M}_G)$ - the weak-star closure of A in \mathcal{M}_G^*
- by the definition of $H^\infty(\mathcal{M}_G)$, the values of its every element are uniquely defined on each $x \in G$

Proposition

$H^\infty(\mathcal{M}_G)$ is isometrically isomorphic to $A^{**} / \mathcal{M}_G^\perp \cap A^{**}$

- G - a Gleason part of A
- $H^\infty(\mathcal{M}_G)$ - the weak-star closure of A in \mathcal{M}_G^*
- by the definition of $H^\infty(\mathcal{M}_G)$, the values of its every element are uniquely defined on each $x \in G$

Proposition

$H^\infty(\mathcal{M}_G)$ is isometrically isomorphic to $A^{**} / \mathcal{M}_G^\perp \cap A^{**}$

Corollary

G is a subset of the spectrum of $H^\infty(\mathcal{M}_G)$

The domination condition for $H^\infty(\mathcal{M}_G)$

$$\|f\| = \sup_{x \in G} |f(x)| \quad \text{for any } f \in H^\infty(\mathcal{M}_G)$$

The domination condition for $H^\infty(\mathcal{M}_G)$

$$\|f\| = \sup_{x \in G} |f(x)| \quad \text{for any } f \in H^\infty(\mathcal{M}_G)$$

Proposition

If $H^\infty(\mathcal{M}_G)$ satisfies the domination condition then the band \mathcal{M}_G is equal to the norm closed linear span of all representing measures for points in G , taken in the quotient space $M(X)/A^\perp$.

The domination condition for $H^\infty(\mathcal{M}_G)$

$$\|f\| = \sup_{x \in G} |f(x)| \quad \text{for any } f \in H^\infty(\mathcal{M}_G)$$

Proposition

If $H^\infty(\mathcal{M}_G)$ satisfies the domination condition then the band \mathcal{M}_G is equal to the norm closed linear span of all representing measures for points in G , taken in the quotient space $M(X)/A^\perp$.

- For $f \in H^\infty(\mathcal{M}_G)$ and $z \in G$ we can define $f(z)$ as the value of f on a representing measure ν_z for z .

The domination condition for $H^\infty(\mathcal{M}_G)$

$$\|f\| = \sup_{x \in G} |f(x)| \quad \text{for any } f \in H^\infty(\mathcal{M}_G)$$

Proposition

If $H^\infty(\mathcal{M}_G)$ satisfies the domination condition then the band \mathcal{M}_G is equal to the norm closed linear span of all representing measures for points in G , taken in the quotient space $M(X)/A^\perp$.

- For $f \in H^\infty(\mathcal{M}_G)$ and $z \in G$ we can define $f(z)$ as the value of f on a representing measure ν_z for z .
- By the weak-star density of A in $H^\infty(\mathcal{M}_G)$, this value $f(z)$ does not depend on the choice of representing measure.

The domination condition for $H^\infty(\mathcal{M}_G)$

$$\|f\| = \sup_{x \in G} |f(x)| \quad \text{for any } f \in H^\infty(\mathcal{M}_G)$$

Proposition

If $H^\infty(\mathcal{M}_G)$ satisfies the domination condition then the band \mathcal{M}_G is equal to the norm closed linear span of all representing measures for points in G , taken in the quotient space $M(X)/A^\perp$.

- For $f \in H^\infty(\mathcal{M}_G)$ and $z \in G$ we can define $f(z)$ as the value of f on a representing measure ν_z for z .
- By the weak-star density of A in $H^\infty(\mathcal{M}_G)$, this value $f(z)$ does not depend on the choice of representing measure.
- So the elements of $H^\infty(\mathcal{M}_G)$ can be regarded as functions on G .

Proposition

If G is a bounded domain in \mathbb{C}^n and $f \in H^\infty(\mathcal{M}_G)$ then the defined above $f(z)$ is a bounded analytic function of $z \in G$.

Proposition

If G is a bounded domain in \mathbb{C}^n and $f \in H^\infty(\mathcal{M}_G)$ then the defined above $f(z)$ is a bounded analytic function of $z \in G$.

Proposition

If G is a star-shaped domain in \mathbb{C}^n such that \overline{G} is the spectrum of $A(\overline{G})$, then the algebras $H^\infty(G)$ and $H^\infty(\mathcal{M}_G)$ are isometrically isomorphic. Hence $H^\infty(G)$ is a dual algebra.

Proposition

If G is a bounded domain in \mathbb{C}^n and $f \in H^\infty(\mathcal{M}_G)$ then the defined above $f(z)$ is a bounded analytic function of $z \in G$.

Proposition

If G is a star-shaped domain in \mathbb{C}^n such that \overline{G} is the spectrum of $A(\overline{G})$, then the algebras $H^\infty(G)$ and $H^\infty(\mathcal{M}_G)$ are isometrically isomorphic. Hence $H^\infty(G)$ is a dual algebra.

Open problem

Is $\sigma(A^{**}) = Y / (A^{**})^\perp$, where Y is the spectrum of $C(X)^{**}$?

Proposition

If G is a bounded domain in \mathbb{C}^n and $f \in H^\infty(\mathcal{M}_G)$ then the defined above $f(z)$ is a bounded analytic function of $z \in G$.

Proposition

If G is a star-shaped domain in \mathbb{C}^n such that \overline{G} is the spectrum of $A(\overline{G})$, then the algebras $H^\infty(G)$ and $H^\infty(\mathcal{M}_G)$ are isometrically isomorphic. Hence $H^\infty(G)$ is a dual algebra.

Open problem

Is $\sigma(A^{**}) = Y / (A^{**})^\perp$, where Y is the spectrum of $C(X)^{**}$?

Consequences

If the above open problem would have a positive solution, then the Corona problem would have a positive solution for the case when $H^\infty(G)$ and $H^\infty(\mathcal{M}_G)$ are isometrically isomorphic.

- $Q = \bigcup_{\alpha} G_{\alpha}$, where G_{α} is a Gleason part of A

- $Q = \bigcup_{\alpha} G_{\alpha}$, where G_{α} is a Gleason part of A
- We say that a measure $\mu \in M(X)$ is an *A-measure* (or analytic measure, or a Henkin measure) with respect to the set Q if $\int u_n d\mu \rightarrow 0$ whenever $\{u_n\}_{n=1}^{\infty} \subset A$ is a bounded sequence converging to 0 pointwise on Q .

- $Q = \bigcup_{\alpha} G_{\alpha}$, where G_{α} is a Gleason part of A
- We say that a measure $\mu \in M(X)$ is an *A-measure* (or analytic measure, or a Henkin measure) with respect to the set Q if $\int u_n d\mu \rightarrow 0$ whenever $\{u_n\}_{n=1}^{\infty} \subset A$ is a bounded sequence converging to 0 pointwise on Q .

A-measures problem for the algebra A at the points of Q

Does the absolute continuity of a measure μ on X with respect to some representing measure of a point $x \in Q$ imply that μ is an A-measure?

- $Q = \bigcup_{\alpha} G_{\alpha}$, where G_{α} is a Gleason part of A
- We say that a measure $\mu \in M(X)$ is an *A-measure* (or analytic measure, or a Henkin measure) with respect to the set Q if $\int u_n d\mu \rightarrow 0$ whenever $\{u_n\}_{n=1}^{\infty} \subset A$ is a bounded sequence converging to 0 pointwise on Q .

A-measures problem for the algebra A at the points of Q

Does the absolute continuity of a measure μ on X with respect to some representing measure of a point $x \in Q$ imply that μ is an A-measure?

Another formulation

Is any measure which is absolutely continuous with respect to an A-measure, itself an A-measure?

Theorem

If G is a Gleason part of A then $H^\infty(\mathcal{M}_G)$ satisfies the domination condition and the A -measures problem for the algebra A at all points of G has a positive solution.

Theorem

If G is a Gleason part of A then $H^\infty(\mathcal{M}_G)$ satisfies the domination condition and the A -measures problem for the algebra A at all points of G has a positive solution.

Theorem

A -measures problem for the algebra A at the points of Q has a positive solution if Q is a union of some Gleason parts of A .

Before our results, A-measures problem was solved positively for two special cases:

Before our results, A-measures problem was solved positively for two special cases:

- by Cole and Range for X being the closure of a strictly pseudoconvex bounded domain Q in \mathbf{C}^n with C^2 boundary, and A being the algebra of all complex continuous functions on X which are holomorphic on its interior Q

Before our results, A-measures problem was solved positively for two special cases:

- by Cole and Range for X being the closure of a strictly pseudoconvex bounded domain Q in \mathbf{C}^n with C^2 boundary, and A being the algebra of all complex continuous functions on X which are holomorphic on its interior Q
- by Bekken and Bui Doan Khanh in the case of the cartesian product of compact planar sets for two classes of algebras - for algebras of continuous functions which are holomorphic on the interior and for algebras generated by rational functions with singularities off X

Before our results, A-measures problem was solved positively for two special cases:

- by Cole and Range for X being the closure of a strictly pseudoconvex bounded domain Q in \mathbf{C}^n with C^2 boundary, and A being the algebra of all complex continuous functions on X which are holomorphic on its interior Q
- by Bekken and Bui Doan Khanh in the case of the cartesian product of compact planar sets for two classes of algebras - for algebras of continuous functions which are holomorphic on the interior and for algebras generated by rational functions with singularities off X

In both above cases the advanced complex analysis methods were used.

THANK YOU!